

Discrete Mathematics 84 (1990) 213–216
North-Holland

213

NOTE

LOWER BOUNDS FOR BI-COLORED QUATERNARY RAMSEY NUMBERS

Aditya SHASTRI

Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

Received 12 January 1988

Revised 16 September 1988

In this note, we prove that $R(5, 5; 4) \geq 19$. We also compute lower bounds for some higher order numbers.

1. Introduction

Let S be a set of cardinality n and if r is a positive integer then $S^{(r)}$ is the family of all r -subsets of S . $S^{(r)} = (R, B)$ denotes partition of $S^{(r)}$ into two classes called “Red” and “Blue”. If $p, q \geq r$ then for sufficiently large n every partition of $S^{(r)}$ into two classes has a p -set, all of whose r -subsets are red (red $K_p^{(r)}$) or a q -set, all of whose r -subsets are blue (blue $K_q^{(r)}$). Minimal such n is the Ramsey number $R(p, q; r)$. A partition without a red $K_p^{(r)}$ and blue $K_q^{(r)}$ is called $(p, q; r)$ -Ramsey partition ($(p, q; r)$ -Ramsey coloring). Though F.P. Ramsey [12] proved the existence of these numbers, their computation is found to be difficult. The most widely studied numbers are bi-colored binary numbers ($r = 2$) (see [2–4, 6–8]). The case $r = 3$ has been studied in [1, 5, 9–11], but only bounds are known even for the first nontrivial case ($13 \leq R(4, 4; 3) \leq 15$). In this note, we compute lower bounds for bi-colored quaternary Ramsey numbers ($R(p, q; 4)$).

2. $R(5, 5; 4)$

Here we give a $(5, 5; 4)$ -Ramsey partition on an 18-set showing $R(5, 5; 4) \geq 19$.

Theorem 1. $R(5, 5; 4) \geq 19$.

Proof. Let F be $GF(17)$, Galois field on 17 elements, and let $S = F \cup \{\infty\}$. We

know that the set of all transformations

$f : S \rightarrow S$ given by

$$f(x) = \begin{cases} \frac{ax+b}{cx+d} & \text{if } ad-bc \neq 0, \quad cx+d \neq 0 \\ \infty & \text{if } x = -\frac{d}{c} \\ \frac{a}{c} & \text{if } x = \infty \end{cases}$$

form a group under composition, call it $L(F)$. Now we partition $S^{(4)}$ into (R, B) by partitioning equivalence classes of $S^{(4)}$ under the induced action of $L(F)$. f in $L(F)$ given by

$$f(x) = \frac{x-b}{x-c} \cdot \frac{a-c}{a-b}$$

maps a, b, c onto $1, 0, \infty$ respectively. Thus, using the above transformation any quadruple $\{a, b, c, d\}$ can be mapped onto $\{1, 0, \infty, d'\}$, which we call a quadruple in standard form (denoted by $\overline{d'}$). The three orbits of $S^{(4)}$ are the following (where only standard quadruples are listed):

$$\begin{aligned} C &= \{\bar{3}, \bar{6}, \bar{8}, \bar{10}, \bar{12}, \bar{15}\}, \\ D &= \{\bar{4}, \bar{5}, \bar{7}, \bar{11}, \bar{13}, \bar{14}\}, \\ E &= \{\bar{2}, \bar{9}, \bar{16}\}. \end{aligned}$$

Now we claim that $B = C \cup E$ and $R = D$ form a $(5, 5; 4)$ -Ramsey partition. To verify the claim, we need to check that every standard quintuple $\{1, 0, \infty, x, y\}$ where $\bar{x}, \bar{y} \in R$ or $\bar{x}, \bar{y} \in B$ have at least one 4-subset on either part. Further, it is enough to check this for quadruples $\{1, 0, \infty, 3, x\}$, $\bar{x} \in B$, $\{1, 0, \infty, 2, y\}$, $\bar{y} \in E$ and $\{1, 0, \infty, 4, z\}$, $\bar{z} \in R$ because there exists a transformation which sends $\{1, 0, \infty, x\}$ to $\{1, 0, \infty, 3\}$ for all $\bar{x} \in C$ and, therefore, $\{1, 0, \infty, x, y\}$ is equivalent to $\{1, 0, \infty, 3, y'\}$. Details are left to the readers. \square

3. Lower bounds for higher order numbers

Abbott and Williams [1] and Kalbfleisch [11] computed lower bounds for some higher order ternary numbers by constructing Ramsey partitions using such partitions for lower order numbers as “building blocks”. We prove

Theorem 2. *If $n \leq R(p, q-1; 4)$ and $m \leq R(p-1, q; 4) - 1$, then $R(p, q; 4) > n + m - 1$.*

Proof. Proof is analogous to a similar theorem of Kalbfleisch [10] for $r = 3$. \square

Table 1

p	q	$n(<R(p, q; 4))$
5	5	18
5	6	24
5	7	29
5	8	36
5	9	44
6	6	46
6	7	75
6	8	111
6	9	154
7	7	4096
7	8	4207
7	9	4361
8	8	8414
8	9	12775
9	9	2^{47}

Corollary. $R(p, q; 4) > 12\binom{p+q-10}{p-5} + \binom{p+q-6}{p-3}$.

Proof. Applying Theorem 2 repeatedly (using induction) and using initial conditions $R(p, 4; 4) = p$ and $R(4, q; 4) = q$, we get

$$R(p, q; 4) > (R(5, 5; 4) - 7) \binom{p+q-10}{p-3} + \binom{p+q-6}{p-3}. \quad \square$$

We must point out, however, that Erdős and Hajnál (see [6], 4.7) found a lower bound for $R(k, k; l)$ for $k \geq l \geq 3$, which in the case of $l = 4$ reduces to $R(k, k; 4) > 2^{2ck^2}$, where c is an absolute constant. Thus, the bounds given above are certainly not the best possible asymptotically. Their bound is based on the so called ‘Stepping-Up Lemma’.

Lemma. If $R(k, k; l) > n$ and $k \geq 3$ then $R(2k + l - 4, 2k + l - 4; l + 1) > 2^n$.

Using Theorem 2, its corollary, Stepping-Up Lemma and lower bounds for $R(4, 4; 3) \geq 13$ [9] and $R(5, 4; 3) \geq 24$ [10], we list lower bounds for some higher order numbers in Table 1.

Acknowledgements

The author wishes to thank Professor Vera T. Sós for her help and encouragement and the referees for their valuable suggestions regarding the bounds in Section 3.

References

- [1] H.L. Abbott and J.R. Williams, Lower bounds for some Ramsey numbers, *J. Combin. Theory Ser. A* 16 (1974) 12–17.
- [2] F.R.K. Chung and C.M. Grinstead, A survey of bounds for classical Ramsey numbers, *J. Graph Theory* 7 (1983) 25–37.
- [3] P. Erdős, Some remarks on the theory of graphs, *Bull. Amer. Math. Soc.* 53 (1947) 292–294.
- [4] P. Erdős, Graph theory and probability II, *Canad. J. Math.* 13 (1961) 346–352.
- [5] G. Giraud, Majoration de nombre de Ramsey ternaire bi-couleur, *C.R. Acad. Sci Paris, Ser. A* 269 (1969) 620–622.
- [6] R. Graham, B. Rothschild and J. Spencer, *Ramsey Theory* (Wiley-Interscience Series, 1980).
- [7] J.E. Graver and J. Yackel, Some graph theoretic results associated with Ramsey's theorem, *J. Combin. Theory* 4 (1968) 125–175.
- [8] R.E. Greenwood and A.M. Gleason, Combinatorial relations and chromatic graphs, *Canad. J. Math.* 7 (1955) 1–7.
- [9] J.R. Isbell, $N(4, 4; 3) \geq 13$, *J. Combin. Theory* 6 (1969) 210.
- [10] J.R. Isbell, $N(5, 4; 3) \geq 24$, *J. Combin. Theory Ser. A* 34 (1983) 379–380.
- [11] J.G. Kalbfleisch, On Robillard's bounds for Ramsey numbers, *Canad. Math. Bull.* 14 (1971) 437–449.
- [12] F.P. Ramsey, On a problem in formal logic, *Proc. London Math. Soc.* 15 (1930) 264–286.